

CABINET:

Daytime Cold & Flu



Active Ingredients (in each softgel)

Purpose

| | |
|---------------------------------|-----------------------------|
| Acetaminophen 325 mg..... | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg..... | Cough suppressant |
| Phenylephrine HCl 5 mg..... | Nasal decongestant |

Uses

temporarily relieves common cold/flu symptoms: • nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches/pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 4 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • thyroid disease • diabetes • high blood pressure • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product • do not use more than directed

Stop use and ask a doctor if • you get nervous, dizzy, or sleepless • symptoms get worse or last more than 5 days (children) or 7 days (adults) • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed - see **Overdose warning** • do not exceed 4 doses per 24 hours • adults & children 12 years of age & over; swallow 2 soft gels with water every 4 hours • children 4 to under 12 years of age; ask a doctor • children under 4 years of age; **do not use**. If taking Daytime and Nighttime soft gels carefully read each label to ensure correct dosing

Other Information

• store at room temperature 15°-30°C(59°-86°F) and avoid excessive heat

Inactive Ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

CABINET:

Nighttime Cold & Flu



Active Ingredients (in each softgel)

Purpose

| | |
|-----------------------------------|-----------------------------|
| Acetaminophen 325 mg..... | Pain reliever/fever reducer |
| Dextromethorphan HBr 15 mg..... | Cough suppressant |
| Doxylamine Succinate 6.25 mg..... | Antihistamine |

Uses

temporarily relieves common cold/flu symptoms: • cough due to minor throat and bronchial irritation • sore throat • headache • minor aches/pains • fever • runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 4 doses in 24 hours, which is the maximum daily amount for this product. • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, lasts for more than 2 days, or occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleep

Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood-thinning drug warfarin

When using this product • do not use more than directed • excitability may occur especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• take only as directed – see **Overdose warning** • do not exceed 4 doses per 24 hours • adults & children 12 years of age & over; swallow 2 soft gels with water every 6 hours • children 4 to under 12 years of age; ask a doctor • children under 4 years of age; do not use. **If taking Daytime and Nighttime products carefully read each label to ensure correct dosing**

Other Information

• store at room temperature 15°-30°C (59°-86°F) and avoid excessive heat • this product does not contain phenylpropanolamine

Inactive Ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

CABINET:

Pain Reliever & Fever Reducer



Active Ingredients (in each tablet)

Purpose

Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Quick medical attention is critical for adults as well as for children

even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- adults & children 12 years & older;
- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor
- children under 12 years; ask a doctor

Other Information

- each tablet contains: magnesium 0.43 mg
- store between 20-25°C (68-77°F)

Inactive Ingredients

Hypermellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, stearic acid, talc, titanium dioxide